

# ECE440 - Introduction to Random Processes

---

## Midterm Exam

November 5, 2014

### Instructions:

- This is an open book, open notes exam.
- Calculators are not needed; laptops, tablets and cell-phones are not allowed.
- Perfect score: 100 (out of 104, extra points are bonus points).
- Duration: 75 minutes.
- This exam has 9 numbered pages, check now that all pages are present.
- Show all your work, and write your final answers in the boxes when provided.

Name: \_\_\_\_\_

Problem	Max. Points	Score	Problem	Max. Points	Score
1.	20		5.	20	
2.	8		6.	16	
3.	8		7.	18	
4.	14				
			Total	104	

**GOOD LUCK!**

1. Suppose that  $X_{\mathbb{N}} = X_0, X_1, \dots, X_n, \dots$  is a Markov chain with state space  $S = \{1, 2\}$  and transition probability matrix

$$\mathbf{P} = \begin{pmatrix} 1/2 & 1/2 \\ 3/4 & 1/4 \end{pmatrix}.$$

To spare you of pointless calculations, if needed you may use that

$$\mathbf{P}^3 = \begin{pmatrix} 19/32 & 13/32 \\ 39/64 & 25/64 \end{pmatrix} = \begin{pmatrix} 0.59 & 0.41 \\ 0.61 & 0.39 \end{pmatrix}.$$

(a) (2 points)  $\mathbf{P}[X_4 = 2 \mid X_3 = 1, X_2 = 2, X_1 = 1] = ?$

(b) (2 points)  $\mathbf{P}[X_5 = 1 \mid X_2 = 1, X_0 = 1] = ?$

(c) (2 points)  $\mathbf{P}[X_3 = 1 \mid X_3 = 2, X_2 = 1, X_1 = 1, X_0 = 2] = ?$

(d) (6 points)  $\mathbb{E} [X_7 \mid X_4 = 2] = ?$

(e) (8 points) Let  $N = \min\{n > 0 : X_n = 2\}$ .  $\mathbb{E} [N \mid X_0 = 1] = ?$

2. (8 points) Consider a probability space  $(S, \mathcal{F}, P[\cdot])$ . Suppose that  $D$  and  $E$  are events in  $\mathcal{F}$  such that  $P[D] = 3/5$  and  $P[E] = 4/5$ . From this information, is it possible to tell if  $D$  and  $E$  are mutually exclusive? Explain.

3. (8 points) Suppose that a random variable  $X$  is Poisson-distributed with parameter  $\lambda > 0$ ; i.e.,  $P[X = x] = e^{-\lambda} \lambda^x / x!$  for  $x = 0, 1, 2, \dots$ . Define the random variable  $Y = qX$ , where  $q$  is a number such that  $0 < q < 1$ . Is  $Y$  Poisson-distributed? Justify your answer.

4. Consider a Markov chain with state space  $S = \{1, 2, 3, 4, 5\}$  and transition probability matrix

$$\mathbf{P} = \begin{pmatrix} 0 & a & b & 0 & 0 \\ c & 0 & d & 0 & 0 \\ e & 0 & 0 & 0 & 0 \\ 0 & 0 & f & 0 & g \\ 0 & 0 & 0 & h & 0 \end{pmatrix}$$

where  $a, b, c, d, e, f, g, h > 0$ .

(a) (8 points) Is the Markov chain irreducible? Explain.

(b) (6 points) Is state 4 transient? Explain.

5. Suppose that  $X_{\mathbb{N}} = X_1, X_2, \dots, X_n, \dots$  is an i.i.d. sequence of random variables, where  $\mathbf{P}[X_1 = 1] = 1/4$ ,  $\mathbf{P}[X_1 = 2] = 1/3$ ,  $\mathbf{P}[X_1 = 3] = 5/12$ .

(a) (10 points) Calculate

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{X_i \geq 2\}$$

and provide justification for the existence of the limit.

(b) (10 points) Specify the distribution of the random variable  $Y$ , defined as

$$Y = \sum_{i=1}^7 \mathbb{I}\{X_i \leq 1\}.$$

6. Suppose that  $X_{\mathbb{N}} = X_0, X_1, \dots, X_n, \dots$  is a Markov chain with state space  $S = \{1, 2\}$  and transition probability matrix

$$\mathbf{P} = \begin{pmatrix} 1/4 & 3/4 \\ 4/5 & 1/5 \end{pmatrix}.$$

(a) (12 points) Compute the stationary distribution of  $X_{\mathbb{N}}$ .

(b) (4 points) Suppose that  $X_0$  has the distribution obtained in part (a).  $\mathbf{P}[X_3 = 1] = ?$

7. (18 points) Suppose that  $X_n$  is the amount of inventory in a store at the end of the time period  $n$ , and that  $D_n$  is the amount of demand that arrives during period  $n$ . Suppose that  $D_{\mathbb{N}} = D_1, D_2, \dots, D_n, \dots$  is an i.i.d. sequence (independent of  $X_0$ ) of non-negative integer-valued random variables, each with probability mass function  $q(\cdot)$ ; i.e.,

$$P[D_1 = i] = q(i), \quad i = 0, 1, 2, \dots$$

At the start of each time period, we receive a shipment of 5 units of inventory to the store. Demand that cannot be met is assumed to go away unsatisfied. Under the preceding assumptions,  $X_{\mathbb{N}} = X_0, X_1, \dots, X_n, \dots$  is a Markov chain with state space  $S = \{0, 1, 2, \dots\}$ . For  $n \geq 0$ , the inventory level at the end of period  $n + 1$  is determined by

$$X_{n+1} = \max\{0, X_n + 5 - D_{n+1}\}.$$

Notice that  $\max\{0, a\} = a$  if  $a \geq 0$ , and  $\max\{0, a\} = 0$  if  $a < 0$ . Hence, the above expression enforces the physical constraint that  $X_{n+1} \geq 0$  always, and  $X_{n+1} = 0$  when the demand  $D_{n+1}$  exceeds the available inventory  $X_n + 5$ .

(a) (6 points) Determine the transition probabilities  $P_{ij}$  for all  $i \geq 0$  and  $j = 1, \dots, i + 5$ .

(b) (6 points) Determine the transition probabilities  $P_{ij}$  for all  $i \geq 0$  and  $j \geq i + 6$ .



(c) (6 points) Determine the transition probabilities  $P_{i0}$  for all  $i \geq 0$ .