

# ECE440 - Introduction to Random Processes

---

## Midterm Exam

October 29, 2018

### Instructions:

- This is an open book, open notes exam.
- Calculators are not needed; laptops, tablets and cell-phones are not allowed.
- Perfect score: 100 points.
- Duration: 90 minutes.
- This exam has 12 numbered pages, check now that all pages are present.
- Make sure you write your name in the space provided below.
- Show all your work, and write your final answers in the boxes when provided.

Name: \_\_\_\_\_

Problem	Max. Points	Score	Problem	Max. Points	Score
1.	28		5.	16	
2.	14		6.	10	
3.	8		7.	16	
4.	8				
			Total	100	

**GOOD LUCK!**

1. Suppose that  $X_{\mathbb{N}} = X_0, X_1, \dots, X_n, \dots$  is a Markov chain with state space  $S = \{1, 2\}$ , transition probability matrix

$$\mathbf{P} = \begin{pmatrix} 1/4 & 3/4 \\ 2/3 & 1/3 \end{pmatrix}$$

and initial distribution  $\mathbf{P}(X_0 = 1) = 1/3$  and  $\mathbf{P}(X_0 = 2) = 2/3$ . To spare you of pointless calculations, if needed you may use that

$$\mathbf{P}^2 = \begin{pmatrix} 9/16 & 7/16 \\ 7/18 & 11/18 \end{pmatrix} = \begin{pmatrix} 0.56 & 0.44 \\ 0.39 & 0.61 \end{pmatrix}.$$

(a) (2 points)  $\mathbf{P}(X_4 = 2 \mid X_2 = 2, X_0 = 1) = ?$

(b) (3 points)  $\mathbf{P}(X_1 = 1, X_0 = 1) = ?$

(c) (4 points)  $P(X_0 = 1 \mid X_1 = 1) = ?$

(d) (4 points)  $\mathbb{E}[X_1] = ?$

(e) (3 points)  $\mathbb{E}[X_2 \mid X_1 = 2] = ?$

(f) (8 points) Compute the stationary distribution of  $X_{\mathbb{N}}$ .

(g) (4 points) Calculate

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=1}^m (X_n)^2$$

and provide justification for the existence of the limit.

2. Consider a random variable  $X$  that is uniformly distributed in the interval  $[0, 1]$ , something we denote as  $X \sim \text{Uniform}[0, 1]$ . Let  $Y$  be a random variable which, conditioned on  $X = x$ , is uniformly distributed over the interval  $[x, 1]$ , that is  $Y | X = x \sim \text{Uniform}[x, 1]$ .

(a) (4 points) What is  $f_{Y|X}(y|x)$ , the conditional probability density function of  $Y$  given  $X = x$ ?

(b) (5 points)  $\mathbb{E}[Y | X = x] = ?$


(c) (5 points)  $\mathbb{E}[Y] = ?$

3. (8 points) Consider a computer program having  $n = 100$  pages of code. Let  $X_i$  be the number of bugs on the  $i$ th page of code. Suppose that the  $X_i, i = 1, \dots, n$ , are i.i.d. random variables having Poisson distribution with mean 1. Let  $Y = \sum_{i=1}^n X_i$  be the total number of bugs.

Use the Central Limit Theorem to approximate  $\mathbf{P}(Y < 90)$ . Write your result in terms of the complementary cumulative distribution function  $\Phi$  of a standard Normal random variable, that is for  $Z \sim \mathcal{N}(0, 1)$  then

$$\Phi(z) = \mathbf{P}(Z \geq z) = \frac{1}{\sqrt{2\pi}} \int_z^{\infty} e^{-u^2/2} du.$$

4. (8 points) The Erdős-Rényi model specifies the simplest mechanism to generate a random graph on  $N$  vertices. It yields undirected graphs (edges do not have directionality) without self loops (an edge connecting a vertex to itself is not allowed). For fixed parameters  $N$  and  $0 \leq p \leq 1$ , the Erdős-Rényi model specifies that each of the possible  $\binom{N}{2}$  edges is included in the graph with probability  $p$ , independently from every other edge. For  $p = 0$ , the graph has no edges. For  $p = 1$ , one obtains a complete graph where every pair of vertices is connected by an edge. A sample realization of an Erdős-Rényi graph with  $N = 20$  and  $p = 0.15$  is shown below.



In graph theory, the degree of a vertex is the number of incident edges to that vertex. In the sample graph above, the degree of vertex 13 is 3 while the degree of vertex 4 is 2. For the Erdős-Rényi model, the degrees  $D_v$  of vertices  $v = 1, \dots, N$  are identically distributed random variables. Name the distribution of the random variable  $D_v$  and specify its parameters.

5. As part of her thesis work, a graduate student from the Warner School is interested in modeling the employment dynamics of young people using a Markov chain. After carrying out a field survey and processing the data, she was able to estimate the following transition probabilities.

	Student	Intern	Employed	Unemployed
Student	0.8	0.1	0.1	0
Intern	0.5	0.5	0	0
Employed	0	0	0.9	0.1
Unemployed	0	0	0.4	0.6

(a) (6 points) Draw the corresponding state transition diagram.

(b) (3 points) Is the Markov chain ergodic? Explain.



(c) (7 points) In the long run, what fraction of time will an individual be unemployed?

6. Consider two random variables  $X$  and  $Y$ . Let  $c$  be a deterministic constant.

(a) (3 points) Derive a simple expression for  $\text{cov}[X, cY]$  in terms of  $c$  and  $\text{cov}[X, Y]$ .

(b) (3 points) Derive a simple expression for  $\text{cov}[X, X + Y]$  in terms of  $\text{var}[X]$  and  $\text{cov}[X, Y]$ .

(c) (4 points) If  $X$  and  $Y$  have a covariance of  $\text{cov}[X, Y]$ , we can transform them to a new pair of random variables whose covariance is zero. To do so, we consider the linear transformation

$$\begin{aligned}W &= X \\Z &= X + aY,\end{aligned}$$

where  $a$  is a deterministic constant. Find the value of  $a$  so that  $W$  and  $Z$  are uncorrelated.

7. During each day, a non-negative integer number of customers arrives to a store to purchase a particular product. Each customer purchases a unit of the product when the product is in stock. Customers who do not find the product in stock depart without making a purchase. The store may order new units of the product at the end of the day (after that day's demand has materialized), and any such orders arrive to the store before the beginning of the next day.

Each day orders are made as follows. If, at the end of the day, there are 5 or fewer units of the product in stock, then an order is placed so that there will be exactly 10 units of inventory present at the start of the next day. If there are more than 5 units of inventory present, no order is placed. Suppose that the daily demand  $D_{\mathbb{N}} = D_1, D_2, \dots, D_n, \dots$  is an i.i.d. sequence of non-negative integer-valued random variables, each with probability mass function  $p(\cdot)$ ; i.e.,

$$P[D_1 = i] = p(i), \quad i = 0, 1, 2, \dots$$

Suppose that at the beginning of the first day of operation ( $n = 0$ ), the stock level is an arbitrary fixed non-negative integer  $z$ .

Let  $X_{\mathbb{N}} = X_0, X_1, \dots, X_n, \dots$  be the Markov chain that represents the amount of the product in stock at the beginning of each day.

(a) (4 points) Determine the transition probabilities  $P_{ij}$  for all  $i \leq 5$  and  $j \geq 0$ .

(b) (4 points) Determine the transition probabilities  $P_{ij}$  for all  $i \geq j > 5$  and  $j \neq 10$ .

(c) (4 points) Determine the transition probabilities  $P_{i10}$  for all  $5 < i < 9$ .

(d) (4 points) Determine the transition probabilities  $P_{i10}$  for all  $i \geq 10$ .